

# ANALOGUE OF SYLVESTER-CAYLEY FORMULA FOR INVARIANTS OF TERNARY FORM

LEONID BEDRATYUK

ABSTRACT. The number  $\nu_d(n)$  of linearly independent homogeneous invariants of degree  $n$  for the ternary form of degree  $d$  is calculated. The following formula is hold

$$\nu_d(n) = c_d(n, 0, 0) + c_d(n, 3, 0) + c_d(n, 0, 3) - 2c_d(n, 1, 1) - c_d(n, 2, 2),$$

here  $c_d(n, i, j)$  is number of nonnegative integer solutions of the system of equations

$$\begin{cases} \sum_i i \alpha_{i,j} = \frac{nd}{3} - \frac{i-j}{3}, \\ \sum_j j \alpha_{i,j} = \frac{nd}{3} - \frac{i+2j}{3}, \\ \sum_{i,j} \alpha_{i,j} = n. \end{cases}$$

Also,  $\nu_d(n)$  is equal to coefficient of  $(pq)^{\frac{nd}{3}}$  in the following polynomial expression

$$(1 + pq + \frac{q^2}{p} - 2q - q^2) \left( \sum_{i_1 + i_2 + \dots + i_d \leq n} \begin{bmatrix} 1 \\ i_1 \end{bmatrix}_{pq} \begin{bmatrix} 2 \\ i_2 \end{bmatrix}_{pq} \dots \begin{bmatrix} d \\ i_d \end{bmatrix}_{pq} \right),$$

here  $\begin{bmatrix} d \\ k \end{bmatrix}_{pq}$  is  $pq$ -binomial coefficient

$$\begin{bmatrix} d \\ k \end{bmatrix}_{pq} := \frac{(p^{d+1} - q^{d+1}) \dots (p^{d+k} - q^{d+k})}{(p - q)(p^2 - q^2) \dots (p^k - q^k)}.$$

1. Let  $T_d$  be the vector  $\mathbb{C}$ -space of ternary forms of degree  $d$  :

$$u(x, y, z) = \sum_{i+j \leq d} \frac{d!}{i!j!(d-(i+j))!} a_{i,j} x^{d-(i+j)} y^i z^j,$$

where  $a_{i,j} \in \mathbb{C}$ . The vector space  $T_d$  is endowed with the natural action of the group  $SL(3, \mathbb{C})$ , and with the corresponding action of the Lie algebra  $\mathfrak{sl}_3$ . Let us identify the algebra of polynomial function  $\mathbb{C}[T_d]$  with the polynomial  $\mathbb{C}$ -algebra  $A_d$  of the variables  $a_{0,0}, a_{1,0}, \dots, a_{0,d}$ . The natural action of the group  $SL(3, \mathbb{C})$  on  $T_d$  induces actions of  $SL(3, \mathbb{C})$  and  $\mathfrak{sl}_3$  on the algebra  $A_d$ . The corresponding invariant ring  $A_d^{SL_3} = A_d^{\mathfrak{sl}_3}$  is called the ring of invariants for the ternary form of degree  $d$ .

The ring  $A_d^{SL_3}$  is graded ring

$$A_d^{SL_3} = (A_d^{SL_3})_0 + (A_d^{SL_3})_1 + \dots + (A_d^{SL_3})_n + \dots,$$

here  $(A_d^{SL_3})_n$  is the vector space generated by homogeneous invariants of degree  $n$ . In this paper we discover the exact formula for calculation the dimension of the vector space  $(R_d^{SL_3})_n$ . The formula is analogue of well-known Sylvester-Cayley formula for invariants of binary form

**2.** To begin with, we give a short proof of Sylvester-Cayley fo invariants of binary form. We will use the idea of the proof in next section to derive an analogous result in the case of ternary form.

Let  $V \cong \mathbb{C}^2$  be standard two-dimensional representation of Lie algebra  $\mathfrak{sl}_2$ . The irreducible representation  $V_d = \langle v_0, v_1, \dots, v_d \rangle$ ,  $\dim V_d = d + 1$  is the symmetric  $d$ -power of the standard representation  $V = V_1$ , i.e.  $V_d = S^d(V)$ ,  $V_0 \cong \mathbb{C}$ . The basis elements  $\hat{D} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ ,  $D = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ ,  $E = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  of the algebra  $\mathfrak{sl}_2$  act on  $V_d$  by the derivations :

$$D(v_i) = i v_{i-1}, \hat{D}(v_i) = (d - i) v_{i+1}, E(v_i) = (d - 2i) v_i.$$

The action of  $\mathfrak{sl}_2$  is extended to action on the symmetrical algebra  $S(V_d)$  in the natural way. The algebra  $I_d$

$$I_d = S(V_d)^{\mathfrak{sl}_2} = \{v \in S(V_d) | D(v) = 0, \hat{D}(v) = 0\},$$

is called the algebra of invariant of the "binary form" of degree  $d$ .

The algebra  $S(V_d)$  is graded

$$S(V_d) = S^0(V_d) + S^1(V_d) + \dots + S^n(V_d) + \dots,$$

and each  $S^n(V_d)$  is complete reducibly respresentation of the Lie algebra  $\mathfrak{sl}_2$ . Thus, the following decomposition is hold

$$S^n(V_d) \cong \gamma_d(n, 0)V_0 + \gamma_d(n, 1)V_1 + \dots + \gamma_d(n, d)V_d, \quad (*)$$

here  $\gamma_d(n, i)$  is the multiplicities of the representation  $V_i$  in the decomposition of  $S^n(V_d)$ . In particular, the multiplicities  $\gamma_d(n, 0)$  of trivial respresentation  $V_0$  is equal to number of linearly independed homogeneous invariants of degree  $n$  for the binary form of degree  $d$ .

The set of weights (the eigenvalues of the operator  $E$ ) of a representatin  $W$  denote by  $\Lambda_W$ , in particular,  $\Lambda_{V_d} = \{-d, -d + 2, \dots, d\}$ .

The formal sum

$$\text{Char}(W) = \sum_{i \in \Lambda_W} n_W(i) q^i,$$

is called the character of the respresentation  $W$ , here  $n_W(i)$  denotes the multiplicities of the weight  $i \in \Lambda_W$ . Since, the multiplicities of any weight of the irreducible respresentation  $V_d$  is equal to 1, we have

$$\text{Char}(V_d) = q^{-d} + q^{-d+2} + \dots + q^d = \frac{q^{d+1} - q^{-(d+1)}}{q - q^{-1}}.$$

The character  $\text{Char}(S^n(V_d))$  of the representation  $S^n(V_d)$  is equal to  $H_d(q^{-d}, q^{-d+2}, \dots, q^d)$ , see [1], where  $H_d(x_0, x_1, \dots, x_d)$  is the complete symmetrical function

$$H_d(x_0, x_1, \dots, x_d) = \sum_{|\alpha|=n} x_0^{\alpha_0} x_1^{\alpha_1} \dots x_d^{\alpha_d}, |\alpha| = \sum_i \alpha_i,$$

see [2] for details.

By putting  $x_i = p^{d-2i}$ ,  $i = 0, \dots, d$ , we obtain the expression for the character of  $\text{Char}(S^n(V_d))$  :

$$\begin{aligned} \text{Char}(S^n(V_d)) &= \sum_{|\alpha|=n} (p^d)^{\alpha_0} (p^{d-2 \cdot 1})^{\alpha_1} \dots (p^{d-2d})^{\alpha_d} = \sum_{|\alpha|=n} p^{dn-2(\alpha_1+2\alpha_2+\dots+d\alpha_d)} = \\ &= \sum_{i=0}^{dn} \omega_d(n, i) p^{dn-2i}, \end{aligned}$$

here  $\omega_d(n, i)$  is the number nonnegative integer solutions of the equation

$$\alpha_1 + 2\alpha_2 + \dots + d\alpha_d = \frac{dn - i}{2}$$

on the assumption that  $|\alpha| = n$ . In particular, the coefficient of  $q^0$  (the multiplicities of zero weight) is equal to  $\omega_d(n, \frac{dn}{2})$ , and the coefficient of  $q^2$  is equal  $\omega_d(n, \frac{dn}{2} - 1)$ .

On the other hand, the decomposition (\*) implies the equality

$$\text{Char}(S^n(V_d)) = \gamma_d(n, 0)\text{Char}(V_0) + \gamma_d(n, 1)\text{Char}(V_1) + \dots + \gamma_d(n, dn)\text{Char}(V_{dn}).$$

Now we can prove

**Theorem 1** (Sylvester-Cayley).

$$\gamma_d(n, 0) = \omega_d(n, \frac{dn}{2}) - \omega_d(n, \frac{dn}{2} - 1).$$

*Proof.* The zero weight occurs once in any irreducible representation  $V_i$ ,  $i = 0, \dots, d$  therefore

$$\omega_d(n, \frac{dn}{2}) = \gamma_d(n, 0) + \gamma_d(n, 1) + \dots + \gamma_d(n, dn).$$

The weight 2 occurs once in any irreducible representation (except trivial)  $V_i$ ,  $i = 1, \dots, d$ , therefore

$$\omega_d(n, \frac{dn}{2} - 1) = \gamma_d(n, 1) + \gamma_d(n, 2) + \dots + \gamma_d(n, dn).$$

Thus

$$\gamma_d(n, 0) = \omega_d(n, \frac{dn}{2}) - \omega_d(n, \frac{dn}{2} - 1).$$

This completes the proof. □

One may show, see [3], that

$$\gamma_d(n, 0) = \left( (1 - q) \left[ \frac{d}{n} \right]_q \right)_{q^{\frac{nd}{2}}},$$

here  $\begin{bmatrix} d \\ n \end{bmatrix}_q$  is the  $q$ -binomial coefficient:

$$\begin{bmatrix} d \\ n \end{bmatrix}_q := \frac{(1 - q^{d+1})(1 - q^{d+2}) \dots (1 - q^{d+n})}{(1 - q)(1 - q^2) \dots (1 - q^n)}.$$

**3.** In the Lie algebra  $\mathfrak{sl}_3$  denote by  $E_{ij}$  the matrix unities. The matrix  $H_1 := E_{11} - E_{22}$  and  $H_2 := E_{22} - E_{33}$  generate the Cartan subalgebra in  $\mathfrak{sl}_3$ . The elements  $H_1, H_2$  act on the  $A_d$ , see [4] for the proof, in the following way

$$H_1(a_{i,j}) = (d - (2i + j))a_{i,j}, H_2(a_{i,j}) = (i - j)a_{i,j}.$$

Every monomial  $a^\alpha := a_{0,0}^{\alpha_{0,0}} a_{1,0}^{\alpha_{1,0}} \dots a_{0,d}^{\alpha_{0,d}}$  of degree  $n$  is the eigenvector of the operators  $H_1, H_2$  with the eigenvalues  $nd - (2\omega_1(\alpha) + \omega_2(\alpha))$  and  $\omega_1(\alpha) - \omega_2(\alpha)$  correspondingly. Here  $\omega_1(\alpha) = \sum_i i \alpha_{i,j}$ ,  $\omega_2(\alpha) = \sum_j j \alpha_{i,j}$ ,  $|\alpha| := \sum_{i,j} \alpha_{i,j} = n$ .

Denote by  $\Gamma_{m_1, m_2}$  the standard irreducible representation of  $\mathfrak{sl}_3$  with the highest weight  $\lambda = (m_1, m_2)$ . It can easily be checked that  $\mathbb{C}^3 \cong \Gamma_{1,0}$ ,  $A_d \cong (S^d(\Gamma_{1,0}))^* \cong \Gamma_{0,d}$ , see details in [1].

Let us recall the definition of the formal character of the representation of the Lie algebra  $\mathfrak{sl}_3$ . Let  $\Lambda$  be a weight lattice of all finitedimension representation of  $\mathfrak{sl}_3$ , and  $\mathbb{Z}(\Lambda)$  be their group ring. The ring  $\mathbb{Z}(\Lambda)$  is free  $\mathbb{Z}$ -module with the basis elements  $e(\lambda)$ ,  $\lambda = (\lambda_1, \lambda_2) \in \Lambda$ ,  $e(\lambda)e(\mu) = e(\lambda + \mu)$ ,  $e(0) = 1$ . Let  $\Lambda_\lambda$  be set of all weights of the representation  $\Gamma_\lambda$ . Then the formal character  $\text{Char}(\Gamma_\lambda)$  is defined, see [5], as formal sum  $\sum_{\mu \in \Lambda_\lambda} n_\lambda(\mu) e(\mu) \in \mathbb{Z}(\Lambda)$ , here  $n_\lambda(\mu)$  is the multiplicities of the weight  $\mu$  in the representation  $\Gamma_\lambda$ . For example, for the highest weight  $\lambda = (1, 1)$  we have

$$\Lambda_{(1,1)} = \{(1, 1), (-1, 2), (1, -2), (0, 0), (-2, 1), (-1, -1)\},$$

therefore

$$\text{Char}(\Gamma_{(1,1)}) = e(1, 1) + e(-1, 2) + e(1, -2) + 2e(0, 0) + e(-2, 1) + e(-1, -1).$$

The basis elements  $a_{i,j}$  of the ring  $A_d$  have weights  $(d - (2i + j), i - j)$ ,  $i + j \leq d$ , of the the multiplicities 1, thus

$$\text{Char}(\Gamma_{0,d}) = \sum_{i+j \leq d} e(d - (2i + j), i - j).$$

The characted of the symmetric power  $S^n(\Gamma_{d,0})$  is the complete symmetric polynomial of degree  $n$  of the variable set  $e(d - (2i + j), i - j)$ ,  $i + j \leq d$ , see [1]. Therefore we have

$$\text{Char}(S^n(\Gamma_{0,d})) = \sum_{|\alpha|=n} e(0, 0)^{\alpha_{0,0}} e(1, 0)^{\alpha_{1,0}} \dots e(0, d)^{\alpha_{0,d}}, |\alpha| = \sum_{i,j} \alpha_{i,j}.$$

After simplification we obtain

$$\text{Char}(S^n(\Gamma_{0,d})) = \sum_{|\alpha|=n} e(nd - 2\omega_1(\alpha) - \omega_2(\alpha), \omega_1(\alpha) - \omega_2(\alpha)) = \sum_{(i,j) \in \Lambda_{(nd,0)}} c_d(n, i, j)(n) e(i, j),$$

here  $c_d(n, i, j)$  is the number of nonnegative integer solutions of the system of equations

$$\begin{cases} 2\omega_1(\alpha) + \omega_2(\alpha) = d n - i, \\ \omega_1(\alpha) - \omega_2(\alpha) = j, \\ |\alpha| = n, \end{cases}$$

or the equivalent system

$$\begin{cases} \omega_1(\alpha) = \frac{d n}{3} - \frac{i - j}{3}, \\ \omega_2(\alpha) = \frac{d n}{3} - \frac{i + 2j}{3}, \\ |\alpha| = n. \end{cases} \quad (**)$$

For  $i = j = 0$  we get  $\omega_1(\alpha) = \omega_2(\alpha)$  and  $n d = 3\omega_1(\alpha)$ . It implies  $n d = 0 \pmod{3}$ . By adding first two equations we obtain  $i - j = 0 \pmod{3}$ . It implies  $i + 2j = 0 \pmod{3}$ .

On the representation  $\Gamma_\lambda$  let us define the value  $E_\lambda$  in the following way

$$E_\lambda = n_\lambda(0, 0) + n_\lambda(3, 0) + n_\lambda(0, 3) - 2n_\lambda(1, 1) - n_\lambda(2, 2).$$

We assume that  $n_\lambda(i, j) = 0$  if  $(i, j) \notin \Lambda_\lambda$ .

The following theorem play crucial role in the calculation.

**Theorem 2.**

$$E_\lambda = \begin{cases} 1, \lambda = (0, 0), \\ 0, \lambda \neq (0, 0). \end{cases}$$

*Proof.* For  $\lambda = (0, 0)$  the statement is obvious.

Suppose now  $\lambda \neq (0, 0)$ . Let us recall that the weight multiplicities of  $\Gamma_{(i,j)}$  increase by one on each of the concentric hexagons of the eigenvalue lattice and are constant on the internal triangles. To prove the theorem let us consider the cases -  $i - j = 0$ ,  $|i - j| > 3$  and  $|i - j| = 3$ .

Let  $\lambda = (m, m)$ ,  $m > 0$ . Then  $n_\lambda(0, 0) = m + 1$ , and it is clear that  $n_\lambda(1, 1) = m$  and  $n_\lambda(3, 0) = n_\lambda(2, 2) = n_\lambda(0, 3) = m - 1$ . It follows that

$$E_\lambda = m + 1 + m - 1 + m - 1 - 2m - (m - 1) = 0.$$

Let  $\lambda = (m, k)$ ,  $|m - k| > 3$ . Then all multiplicities  $n_\lambda(0, 0)$ ,  $n_\lambda(3, 0)$ ,  $n_\lambda(1, 1)$ ,  $n_\lambda(2, 2)$ ,  $n_\lambda(0, 3)$  are equal to  $\min(m, k) + 1$  and therefore  $E_\lambda = 0$ .

Let  $\lambda = (m, k)$ ,  $m - k = 3$ . Then  $n_\lambda(0, 0) = n_\lambda(3, 0) = n_\lambda(1, 1) = k + 1$  and  $n_\lambda(2, 2) = k$ ,  $n_\lambda(0, 3) = k$ ,  $E_\lambda = 0$ .

If  $\lambda = (m, k)$ ,  $k - m = 3$  then  $n_\lambda(0, 0) = n_\lambda(0, 3) = n_\lambda(1, 1) = m + 1$  and  $n_\lambda(2, 2) = m$ ,  $n_\lambda(3, 0) = m$ . This yields that  $E_\lambda = 0$ .  $\square$

Now we are ready to calculate the value  $\nu_d(n)$ .

**Theorem 3.** *The number  $\nu_d(n)$  of linearly independent homogeneous invariants of degree  $n$  for the ternary form of degree  $d$  is equal*

$$\nu_d(n) = c_d(n, 0, 0) + c_d(n, 3, 0) + c_d(n, 0, 3) - 2c_d(n, 1, 1) - c_d(n, 2, 2).$$

*Proof.* The number  $\nu_d(n)$  is equal to the multiplicities  $\gamma_d(0, 0)$  of trivial representation  $\Gamma_{0,0}$  in the symmetrical power  $S^n(\Gamma_{0,d})$ . Suppose the decomposition is hold

$$S^n(\Gamma_{0,d}) = \gamma_d(0, 0)\Gamma_{0,0} + \cdots + \gamma_d(0, n\,d)\Gamma_{0,n\,d} = \sum_{\lambda \in \Lambda_{(0,n\,d)}^+} \Gamma_\lambda.$$

It implies,

$$\text{Char}(S^n(\Gamma_{0,d})) = \gamma_d(0, 0)\text{Char}(\Gamma_{0,0}) + \cdots + \gamma_d(0, n\,d)\text{Char}(\Gamma_{0,n\,d}).$$

Thus,

$$\sum_{(i,j) \in \Lambda_{(0,n\,d)}} c_d(n, i, j)(n)e(i, j) = \sum_{\lambda} \gamma_d(\lambda)\text{Char}(\Gamma_\lambda) = \sum_{(i,j) \in \Lambda_{(0,n\,d)}} \sum_{\lambda} \gamma_d(\lambda)n_\lambda(i, j)e(i, j).$$

It follows that  $c_d(n, i, j) = \sum_{\lambda} \gamma_d(\lambda)n_\lambda(i, j)$ . By using previous theorem we have

$$c_n(n, 0, 0) + c_d(n, 3, 0) + c_d(n, 0, 3) - 2c_d(n, 1, 1) - c_d(n, 2, 2) = \sum_{\lambda} \gamma_d(\lambda)E_\lambda = \gamma_d(0, 0).$$

Taking into account  $\gamma_d(0, 0) = \nu_d(n)$  we obtain the statement of the theorem.  $\square$

**4.** Let us derive the fourmula for calculation of  $\nu_d(n)$ .

It is not hard to prove that the number  $c_d(n, 0, 0)$  of nonnegative integer solutions of the following system

$$\begin{cases} \omega_1(\alpha) = \frac{dn}{3}, \\ \omega_2(\alpha) = \frac{dn}{3}, \\ |\alpha| = n \end{cases}$$

is equal to coefficient of  $t^n(pq)^{\frac{dn}{3}}$  the expansion of the following series

$$R_d = \left( \prod_{k+l \leq d} (1 - tp^k q^l) \right)^{-1}.$$

Denote it in such way  $c_d(n, 0, 0) = (R_d)_{t^n(pq)^{\frac{dn}{3}}}$ .

The number  $c_d(n, 3, 0)$  of nonnegative integer solutions of the following system

$$\begin{cases} \omega_1(\alpha) = \frac{dn}{3} - 1, \\ \omega_2(\alpha) = \frac{dn}{3} - 1, \\ |\alpha| = n \end{cases}$$

is equal to  $(R_d)_{t^n(pq)^{\frac{dn}{3}-1}} = (pqR_d)_{t^n(pq)^{\frac{dn}{3}}}$ . Similarly,

$$c_d(n, 0, 3) = \left( \frac{q^2}{p} R_d \right)_{t^n(pq)^{\frac{dn}{3}}}, c_d(n, 1, 1) = (qR_d)_{t^n(pq)^{\frac{dn}{3}}}, c_d(n, 2, 2) = (q^2 R_d)_{t^n(pq)^{\frac{dn}{3}}}.$$

Therefore the following formula is hold

$$\nu_d(n) = \left( \left( 1 + pq + \frac{q^2}{p} - 2q - q^2 \right) R_d \right)_{t^n(pq)^{\frac{dn}{3}}}.$$

Let us define the  $pq$ -binomial coefficients in the following way

$$\left[ \begin{matrix} n \\ k \end{matrix} \right]_{pq} := \frac{(p^{n+1} - q^{n+1}) \dots (p^{n+k} - q^{n+k})}{(p - q)(p^2 - q^2) \dots (p^k - q^k)}.$$

The function  $G_m := (\prod_{k+l=m} (1 - tp^k q^l))^{-1}$  are generating functions for the  $pq$ -binomial coefficients:

$$G_m = \prod_{k+l=m} ((1 - tp^k q^l)^{-1}) = 1 + \left[ \begin{matrix} m \\ 1 \end{matrix} \right]_{pq} t + \left[ \begin{matrix} m \\ 2 \end{matrix} \right]_{pq} t^2 + \dots$$

Therefore,

$$R_d = G_0 G_1 \dots G_d = \sum_n \left( \sum_{i_1+i_2+\dots+i_d \leq k} \binom{1}{i_1}_{pq} \binom{2}{i_2}_{pq} \dots \binom{d}{i_d}_{pq} \right) t^n.$$

Thus, the number  $\nu_d(n)$  of linearly independent homogeneous invariants of degree  $n$  for the ternary form of degree  $d$  is calculating by the formula

$$\nu_d(n) = \left( (1 + pq + \frac{q^2}{p} - 2q - q^2) \sum_{i_1+i_2+\dots+i_d \leq k} \binom{1}{i_1}_{pq} \binom{2}{i_2}_{pq} \dots \binom{d}{i_d}_{pq} \right)_{(pq)^{\frac{d}{3}}}.$$

Below are present the first several terms of Poincaré series for the rings of invariants of the ternary forms of degrees 3,4,5,6,7.

$$\begin{aligned} P_3(t) &= t^4 + t^6 + t^8 + t^{10} + 2t^{12} + t^{14} + 2t^{16} + 2t^{18} + 2t^{20} + 2t^{22} + 3t^{24} + 2t^{26} + \dots \\ P_4(t) &= t^3 + 2t^6 + 4t^9 + 7t^{12} + 11t^{15} + 19t^{18} + 29t^{21} + 44t^{24} + 67t^{27} + 98t^{30} + \dots \\ P_5(t) &= 2t^6 + t^9 + 19t^{12} + 24t^{15} + 178t^{18} + 383t^{21} + 1470t^{24} + 3331t^{27} + 9381t^{30} + \dots \\ P_6(t) &= t^3 + t^4 + t^5 + 4t^6 + 5t^7 + 8t^8 + 17t^9 + 28t^{10} + 48t^{11} + 99t^{12} + 172t^{13} + \dots \\ P_7(t) &= 3t^6 + 13t^9 + 421t^{12} + 4992t^{15} + 60303t^{18} + 548966t^{21} + \dots \end{aligned}$$

The results in the case  $d = 4$  coincides completely with the results of Shioda, see [6].

## REFERENCES

- [1] W.Fulton, J. Harris, Representation theory: a first course, 1991.
- [2] I. G. Macdonald, Symmetric Functions and Hall Polynomials, Clarendon Press, Oxford, 1979.
- [3] D. Hilbert, Theory of algebraic invariants. Lectures. Cambridge University Press, (1993).
- [4] L.P. Bedratyuk, *Analogue of Roberts' theorem for the ternary forms*, Nauk. Visn. Chernivets'kogo Univ., Mat., (2007)
- [5] J. Humphreys, Introduction to Lie Algebras and Representation Theory, 1978.
- [6] Shioda, T., On the graded ring of invariants of binary octavics. *Am. J. Math.*, 89, 1022–1046, (1967).

KHMEL'NYTS'KY NATIONAL UNIVERSITY, INSTYTUTS'KA ST. 11, KHMEL'NYTS'KY, 29016, UKRAINE  
E-mail address: bedratyuk@ief.tup.km.ua